

Determination of Stability Constant of Trimethoprim-Zn(II) complex at Different Temperatures by Continuous Variation Method

O.V. Ikpeazu¹, I.E. Otuokere^{2*}, K.K.Igwe³

¹Department of Biochemistry, Abia State University, Uturu, Nigeria

^{2*}Department of Chemistry, Michael Okpara University of Agriculture, Umudike, Nigeria

³Department of Vet. Biochemistry and Animal Production, Michael Okpara University of Agriculture, Umudike, Nigeria

Corresponding authors' E.mail: ifeanyiotuokere@gmail.com

Abstract: Trimethoprim is an antibiotic that is primarily used in the treatment of urinary tract infections. Continuous variation method was applied in the calculation of stability constant of trimethoprim-Zn(II) complex depending on the theoretical explanation of the stoichiometry. The formation of Zn(II) complex with trimethoprim was studied colorimetrically at an absorption maximum of 630 nm at 25, 30, 35 and 40 °C. The data showed that Zn(II) and trimethoprim combine in the molar ratio of 1:1 at pH 7.4 with ionic strength maintained using 0.1M KNO₃. Calculated stability constants values were 8.13×10^6 , 9.21×10^5 , 4.47×10^5 and 4.52×10^4 at 25, 30, 35 and 40 °C respectively. Calculated ΔG° for the complex were -3.94×10^4 , -3.46×10^4 , -3.33×10^4 and -2.79×10^4 at 25, 30, 35 and 40 °C respectively. The stability constant results suggested that trimethoprim used in the study is a potent chelating agent and can be an efficient antidote in the therapy of Zn(II) overload or poisoning.

Keywords: Absorbance, colorimetric, complex, stability constant, trimethoprim, zinc.

I. INTRODUCTION

The human body has an elaborate mechanism for management and regulation of key trace metals circulating in bloodstream and in the cells. Essential trace metals from our diet are transported into the blood when there is depletion in blood levels. Essential trace metals are also transported into cells if there is inadequacy in the cellular levels. Excreted of trace metals occur if blood and cellular levels are sufficiently overloaded. When there is systemic failure in function, abnormality in levels and ratios of trace metals can develop. One of such trace-metal imbalances is depressed zinc. The concentration of zinc is important for healthy living [1]. The amount of Zn distributed in the human body is about 2-4 grams [2]. Zinc is found mainly in the brain, muscle, bones, kidney and liver, with the highest concentrations in the prostate and parts of the eye [3]. Zn is the second most dominant transition metal in biological systems after iron and it is the only metal that appears in all enzymatic classes [2,4]. Many cellular metabolic processes requires the involvement of Zn [5]. It was reported that 10% of human proteins potentially complex with zinc, including hundreds which transport and traffic zinc [6,7]. Catalytic activity of more than 200 enzymes requires Zn mineral [6,7]. It is essential in boosting immunity [7,8], facilitate wound healing [8] and the synthesis of protein.

Trimethoprim (Figure 1) is an antibiotic that is primarily used in the treatment of urinary tract infections [9]. Other uses include for middle ear infections and travelers' diarrhea [9]. With sulfamethoxazole or dapsone it may be used for *Pneumocystis* pneumonia in people with HIV/AIDS [9] It is administered orally [9]. Common side effects include nausea, changes in taste, and rash [9]. Rarely it may result in blood problems such as not enough platelets or white blood cells [9].




Figure 1: The chemical structure of trimethoprim

The Synthesis and characterization of trimethoprim metal complexes (Figure 2) have been reported [10]. Trimethoprim metal complexes were characterized by different spectroscopic methods: Ultraviolet-visible, Fourier transform infrared, conductivity measurements, thermal analysis (TG) and magnetic susceptibility measurement [10]. The results of the experiment showed that the coordination of trimethoprim drug with the transition metal ions occurred through nitrogen of pyrimidinyl ring. Square planar geometry was suggested for Mn(II), Ni(II), Cu(II) complexes while Cr(III) and Co(II) complexes have an octahedral geometry. The complexes have electrolyte properties [10].




Figure 2: Square planar geometry of trimethoprim metal complex [10].

For several decades, chelating agents have been used as antidote to combat metal poisoning [11, 12]. Biological friendly sequestering agents have been used effectively to chelate metals in patients with metal overload [11-13]. However, chelating capacity is a function of stability constant indicating that the effectiveness of a drug to chelate with a metal ion depends on the stability constant and other parameters [11-13]. Many authors have reported the study of stability constant of drug- metal complexes [13-16]. However, to the best of our knowledge, the stability constant of trimethoprim-Zn(II) complex at different temperatures have not been reported elsewhere in literature. Therefore, the present study is aimed at determining the stability constant of trimethoprim-Zn(II) complex using continuous variation. Information on stability constants of this complex can be useful in analysing the effects of trimethoprim on zinc ion and other electroactive divalent trace metals. It is possible that changes in trace metal and mineral concentration induced by trimethoprim can be an efficient antidote in the therapy of Zn overload or poisoning.

II. MATERIALS AND METHODS

Reagents

Reagents used for the study were of analytical grade. Trimethoprim was purchased from Andhra Organics Limited, Indian. ZnCl₂ was purchased from Merck & Co., Inc USA. Double-distilled water was used throughout the experiment.

Preparation of 1 x 10⁻² M ZnCl₂

ZnCl₂ (1.362 g, 10 mmol, M. Wt. = 136.290 g/mol) was dissolved in freshly distilled water in a 250 cm³ beaker and was made up to the mark in a 1000 cm³ volumetric flask.

Preparation of 1 x 10⁻² M trimethoprim

Trimethoprim (2.903 g, 10 mmol, molar weight = 290.32 g/mol) was dissolved in freshly distilled water in a 250 cm³ beaker and was made up to the mark in a 1000 cm³ volumetric flask.

Procedure for continuous variation method

Exactly 0, 1, 2, 3, 4, 5, 6 cm³ of 1 x 10⁻² M ZnCl₂ were pipetted into seven different 50 cm³ volumetric flasks respectively. Exactly 6, 5, 4, 3, 2, 1, 0 cm³ of 1 x 10⁻² M of trimethoprim was added to the respective flasks containing Zn(II) solution. The pH was adjusted to 7.4 while the ionic strength was maintained constant using 0.1 M KNO₃. The absorbance of each solution was measured at 630 nm (maximum wavelength of absorbance of the complex) and at temperatures of 25 and 40 °C, respectively.

Calculation of stoichiometry, stability constant and free energy

The stoichiometry mole fraction (SMF) of the complex using continuous variation method was calculated using equation 1 [17].

$$SMF = \frac{m}{1-m} \tag{1}$$

where m is the mole fraction of the metal ion. The stability constant was calculated using the classical method expressed in equation 2,

$$K_{st} = \frac{1-\alpha}{m^m \cdot n^n (\alpha)^{m+n} (C)^{m+n-1}} \tag{2}$$

where C is the concentration of the complex at stoichiometry point, α is the degree of dissociation, m and n are the corresponding stoichiometric coefficients of metal and ligand respectively. The degree of dissociation (α) was calculated using equations 3, 4 and 5 [17].

$$A_{\alpha} = A_o - A_{max} \tag{3}$$

$$A_{max} = \epsilon b C \tag{4}$$

$$\alpha = \frac{A_{\alpha}}{\epsilon b C} \tag{5}$$

where A_{max} is absorbance value of the maximum at experimental curve that represents the maximum quantity of the complex that is formed. A_o is absorbance value corresponding to the intersect point of the theoretical straight lines. A_α is the absorbance value of the part of dissociated concentration of complex. ε is molar absorptivity, b is cell thickness, C is a concentration of complex at stoichiometry point.

The Gibbs free energy was calculated using the Helmholtz Gibb equation (equation 6),

$$\Delta G^{\theta} = -RT \ln K \tag{6}$$

III. RESULTS AND DISCUSSION




Fig. 3: Absorption spectra of ZnCl₂ (1 x 10⁻² M) (series 1) and trimethoprim-Zn(II) complex (series 2)

The absorption spectra (Fig. 3) shows the absorbance of $ZnCl_2$ (series 1) and trimethoprim-Zn(II) complex (series 2) at wavelength of 400 – 700 nm. The absorption spectra showed that the wavelength of maximum absorbance of the complex was 630 nm. At this wavelength, $ZnCl_2$ displayed minimal absorbance. Since the maximum absorbance of the complex was 480 nm, it was used for the analytical measurement in the determination of the stability constants and free energies. The maximum absorbance of $ZnCl_2$ was observed at wavelength of 520 nm. It was observed that trimethoprim-Zn(II) complex gave a water soluble complex in aqueous solution, This may be attributed to the ability of water to act as a weak monodentate ligand in forming labile Zn-aquo complex. During complexation, trimethoprim displaced water from Zn-aquo to form a stable trimethoprim-Zn(II) complex. Similar labile aquo complexes were also reported by several authors in their study of stability constant of complexes [11-17].

Table I: Experimental data of trimethoprim-Zn(II) complex at 630 nm by continuous variation method

S/N	ZnCl ₂ (1x 10 ⁻² M)	Trimethoprim (1 x 10 ⁻² M)	Mole fraction of Zn(II)	Absorbance at 630 (nm)			
				25 °C	30 °C	35 °C	40 °C
1	0.000	6.000	0.000	0.00	0.00	0.00	0.00
2	1.000	5.000	0.170	0.00	0.00	0.00	0.00
3	2.000	4.000	0.330	0.00	0.00	0.01	0.00
4	3.000	3.000	0.500	0.01	0.02	0.02	0.02
5	4.000	2.000	0.660	0.03	0.03	0.03	0.03
6	5.000	1.000	0.830	0.01	0.01	0.02	0.02
7	6.000	0.000	1.000	0.00	0.00	0.00	0.00




Figure 4: Job's curve of equimolar solutions at 25 °C




Figure 5: Job's curve of equimolar solutions at 30 °C




Figure 6: Job's curve of equimolar solutions at 35 °C




Figure 7: Job's curve of equimolar solutions at 40 °C

The Job's curves at 25, 30, 35 and 40 °C are shown in Figures 4, 5, 6 and 7 respectively. Equation 1 was applied in calculation of stoichiometry of the complex

$$SMF = \frac{0.66}{0.34} = 1.94 \approx 2 \text{ (25 } ^\circ\text{C)}, SMF = \frac{0.66}{0.34} = 1.94 \approx 2 \text{ (30 } ^\circ\text{C)}, SMF = \frac{0.66}{0.34} = 1.94 \approx 2 \text{ (35 } ^\circ\text{C)}, SMF = \frac{0.70}{0.30} = 2.33 \approx 2 \text{ (40 } ^\circ\text{C)}.$$

The stoichiometry calculation corresponded to metal:ligand ratio of 1:2. The extrapolated value at the point of cross-section on continuous variation plot (Figs. 4, 5, 6 and 7) corresponded to the total absorbance of the complex, indicating that the complex formation process has been completed. Several authors have also applied continuous variation method in the determination of metal:ligand ratio in complexes [11-18].

Table II: Calculated stability constant and Gibbs free energies of trimethoprim-Zn(II) complex using continuous variation method

S/N	Temperature (°C)	M:L ratio	Stability constant	ΔG^\ominus (J)
1	25	1:2	8.13×10^6	-3.94×10^4
2	30	1:2	9.21×10^5	-3.46×10^4
3	35	1:2	4.47×10^5	-3.33×10^4
4	40	1:2	4.52×10^4	-2.79×10^4

Calculation of the stability constant and Gibbs free energies were based on equations 2, 3, 4, 5 and 6 respectively. The large values of the stability constant indicate that the metal has high affinity for the ligand, provided the system is at equilibrium. Stability constant is an evaluation of the strength of the interaction between the reagents that come together to form the complex. The values of the stability constant showed that the complex was stable at 25, 30, 35 and 40 °C. Increasing the temperature of coordination from 25 to 40 °C decrease the stability constant values. This suggested that the complex formation reaction is exothermic reaction. The values of the stability constants were positive, indicating that the complex is stable. Positive stability constant values using continuous variation have also been reported by several authors [11-18]. Continuous variation is an established techniques in the determination of stability constant and Gibbs free energies. The results of stability constant suggested that trimethoprim could be effective in chelation therapy against Zn(II) toxicity. The negative values of the free energies suggested that the complexes were formed spontaneously.

IV. CONCLUSION

The Job's continuous variation methods data showed that Zn(II) and trimethoprim combine in the molar ratio of 1:2. The stability constant results suggested that the complex was stable. Trimethoprim used in the study is a potent chelating agent and can be an efficient antidote in the therapy of Zn(II) overload or poisoning.

REFERENCES

- [1] Kaslow JE. Copper/Zinc Imbalance. Medical Board of California.
- [2] R.A. Wapnir, "Protein Nutrition and Mineral Absorption", CRC Press. 1990
- [3] C.C. Pfeiffer & E.R. Braverman. "Zinc, the brain and behavior", Biol Psychiatry, Vol. 17, pp. 513, 1982
- [4] M.R. Broadley, P.J. White, J.P. Hammond, I. Zelko & A. Lux, "Zinc in plants. New Phytologist", Vol. 173, pp. 677, 2007
- [5] H.G. Classen, U. Gröber, D. Löw, J. Schmidt & H. Stracke "Zinc deficiency: Symptoms, causes, diagnosis and therapy. Med Monatsschr Pharm, Vol. 34, pp. 87, 2001
- [6] H.H. Sandstead, "Understanding zinc: recent observations and interpretations", J Lab Clin Med, Vol. 124, pp. 322, 1994
- [7] T.J. McCarthy, J.J. Zeelie & D.J. Krause, "The antimicrobial action of zinc ion/antioxidant combinations", Clinical Pharmacology & Therapeutics Vol. 17, pp. 5, 1992
- [8] N.W. Solomons, "Mild human zinc deficiency produces an imbalance between cell-mediated and humoral immunity", Nutr Rev, Vol. 56, pp. 27, 1998
- [9] "Trimethoprim". *The American Society of Health-System Pharmacists. Archived from the original on 2015-09-24. Retrieved Aug 1, 2015.*
- [10] F.M. Ibrahim, R.A. Hammza & D.H. Fadhil, Synthesis and characterization of Trimethoprim metal complexes used as corrosion inhibitors for carbon steel in acid media, Int. J. Corros. Scale Inhib., Vol. 8, No. 3, pp. 733–742, 2019
- [11] Ikpeazu, O.V., Otuokere, I.E. and Igwe, K.K (2020) Acetaminophen-Mn(II) complex: determination of stoichiometry, stability constants and Gibbs free energies at different temperatures, Nigerian Research Journal of Chemical Sciences, 8(2): 98 -110
- [12] O. V. Ikpeazu, I. E. Otuokere & K. K. Igwe, Determination of Stability Constants and Gibbs Free Energies of Cefotaxime-Zn(II) Complex at Different Temperatures, International Journal of Trend in Scientific Research and Development, Vol. 4, No.5, pp 166 – 171, 2020
- [13] A.C. Tella, & J.A. Obaleye, "Metal-Chelator Therapy: Stability Constant of Transition metal complexes of Pyrimidine and Sulphonamide Drugs. Int. Chem. Sci. Vol. 8, No. 3, pp. 1675, 2010
- [14] S.A. Tirmizi, M.H. Sarwar, W.S. Sarwar & W. Anwar, "Spectrophotometric study of stability constants of famotidine-Cu(II) complex at different temperatures", The Arabian Journal for Science and Engineering, Vol. 34, No. (2A), pp. 43, 2008

- [15] S.A. Tirmizi, F.H. Wattoo, M.H.S Wattoo, S. Sarwar, A.N. Memon, & A.B. Ghangro “Spectrophotometric study of stability constants of cimetidine–Ni(II) complex at different temperatures”, *Arabian Journal of Chemistry*, Vol. 5, pp. 309 – 314. 2012
- [16] R. Waranyoupalin, R. Wongnawa, S. Wongnawa, M.C. Pakawatchai, M C. Panichayupakaranant, & P. Sherdshoopongse, “Studies of complex formation between curcumin and Hg(II) by spectrophotometric method: A new approach to overcome peak overlap”, *Cent. Eur. J. Chem.*, Vol. 7, No. 3, pp. 388, 2009
- [17] R.F. Abbas, Spectrophotometric determination of stability constant by classical and Modified Varagas equations for procaine penicillin G using diazotization reaction depending on stoichiometric curves, *International Journal of ChemTech Research*, Vol. 10, No 2, pp. 485 – 496, 2017
- [18] O.V. Ikpeazu, I.E. Otuokere & K.K Igwe, Colorimetric Determination of Stability Constant of Acetaminophen-Cu(II) complex by Classical Equation Depending on Stoichiometric Curves, *Communications in Physical Sciences*, Vol. 5, No. 3, pp. 371 – 377. 2020